Light-front quantization

From the White Paper by the Board of Directors of ILCAC, Inc.

John R. Hiller


jhiller@d.umn.edu

Department of Physics
University of Minnesota Duluth



Light-front Quantization

- ILCAC, Inc. URL: http://www.ilcacinc.org/
- framework for analysis of hadrons in QCD: $x^+ = t + z$, $x^- = t z$, $p^- = E p_z$, $p^+ = E + p_z$



nonperturbative Hamiltonian eigenvalue problem:

$$\mathcal{P}^-|P^+,\vec{P}_\perp\rangle = \frac{M^2 + P_\perp^2}{P^+}|P^+,\vec{P}_\perp\rangle$$


• well-defined Fock-state expansion: $|\operatorname{proton}\rangle = \psi_{uud}|uud\rangle + \psi_{uudq}|uudg\rangle + \psi_{uudq\bar{q}}|uudq\bar{q}\rangle + \cdots$


- boost-invariant & process-independent wave functions
 - free of vacuum contributions
- Minkowski space: time-like and space-like processes
- no fermion doubling
- no formulation-specific light-quark problem




Progress & Successes

- direct evidence for a Maldacena duality conjecture
 SDLCQ Collaboration (Pinsky, jrh, et al.)
- massively parallel Lanczos code Vary et al.
- nonperturbative QED in an arbitrary covariant gauge
 Chabysheva and jrh
- relativistic quark models based on light-front holography
 Brodsky and De Téramond
- effective-particle representation via renormalization gp
 Wilson and Głazek
- methods for finite-temperature calculations
 - -SDLCQ Collaboration, Beyer and Strauss









Goals & Challenges

- calculation of hadron masses, wave functions, form factors, GPDs,...
 - light-front coupled-cluster method



- at zero and finite temperature
- understanding of vacuum structure
 - zero modes, effective interactions
- construction of nonperturbative regularization
 - St Petersburg group (Paston et al.)
- classification of total angular momentum eigenstates
 - rotations about x and y are dynamical
- physics of intense laser fields
 - nonperturbative strong-field QED

